## Shieldex Weed Control and Crop Safety Janesville, WI 2020

#### **Trial Information**

Trial Sponsor:	Summit Agro USA
Objective:	Compare Shieldex applied POST to industry standards for weed control and crop safety
Crop:	Corn
Hybrid:	Dekalb DKC 54-38 RIB
Weed species (pressure):	Giant ragweed (high-very high)
Herbicide Application(s):	PRE – 4/28
	POST – 5/27 (V2 corn)

For the full report see trial #20-ROK-CN04 in the 2020 Wisconsin Weed Science Research Report



# Shieldex Weed Control and Crop Safety Janesville, WI 2020

	ΑΡΡ	Giant Ragweed Control %		Corn Injury* %	
Herbicide Treatment*	Timing	6/3 <sup>B</sup>	6/10 <sup>T</sup>	6/3	6/10
1.67 qt Bicep II Magnum	PRE	55	48	0	0
1.67 qt Bicep II Magnum <b>fb</b> 1 fl oz Shieldex + 1 pt atrazine + 1% v/v CO	PRE C POST	100	78	10	5.5
1.67 qt Bicep II Magnum <b>fb</b> 1 fl oz Impact + 1 pt atrazine + 1% v/v COC	PRE POST	99	78	2	0
1.67 qt Bicep II Magnum <b>fb</b> 3 fl oz Laudis + 1 pt atrazine + 1% v/v COC	PRE POST	100	83	1	0
Untreated Check	-	0	0	0	0
	LSD (0.10)	3	9	0.8	0.5
	p-value	<0.01	<0.01	<0.01	<0.01

\*Corn injury rated as % leaf necrosis (leaf burn). Injury was not severe enough to affect normal growth and new leaves were unaffected.

<sup>B</sup> Rated % burndown control

<sup>T</sup>Rated % total control

## Shieldex Weed Control and Crop Safety Janesville, WI 2020

Plot pictures taken on 5/27 29 days after the PRE application The day of the POST application

**Untreated Check** 




PRE: 1.67 qt Bicep II Magnum

## Shieldex Weed Control and Crop Safety Janesville, WI 2020

### Corn Injury - Necrosis

1 fl oz Shieldex + 1 pt atrazine 4L + 1% v/v COC



Wisconsin Cropping Systems Weed Science

©Dr. Rodrigo Werle, University of Wisconsin-Madison

## Plot pictures taken on 6/3 36 days after the PRE application 7 days after the POST application

The number in the upper right-hand corner is the average % burndown control of giant ragweed of 4 replications. Newly emerged giant ragweed seedlings seen in the pictures were not rated at this timing.

#### 36 days after PRE (6/3) Untreated Check



#### 36 days after PRE (6/3) PRE: 1.67 qt Bicep II Magnum



©Dr. Rodrigo Werle, University of Wisconsin-Madison

#### 7 days after POST (6/3) PRE: 1.67 qt Bicep II Magnum *fb* POST: 1 fl oz Shieldex + 1 pt atrazine 4L + 1% v/v COC



©Dr. Rodrigo Werle, University of Wisconsin-Madison

#### 7 days after POST (6/3) PRE: 1.67 qt Bicep II Magnum *fb* POST: 1 fl oz Impact + 1 pt atrazine 4L + 1% v/v COC



©Dr. Rodrigo Werle, University of Wisconsin-Madison

#### 7 days after POST (6/3) PRE: 1.67 qt Bicep II Magnum *fb* POST: 3 fl oz Laudis + 1 pt atrazine 4L + 1% v/v COC



©Dr. Rodrigo Werle, University of Wisconsin-Madison

## Plot pictures taken on 6/10 43 days after the PRE application 14 days after the POST application

The number in the upper right-hand corner is the average % control of giant ragweed of 4 replications. Giant ragweed seedlings that emerged after application were included in this rating.

#### 43 days after PRE (6/10) Untreated Check



#### 43 days after PRE (6/10) PRE: 1.67 qt Bicep II Magnum



#### 14 days after POST (6/10) PRE: 1.67 qt Bicep II Magnum *fb* POST: 1 fl oz Shieldex + 1 pt atrazine 4L + 1% v/v COC



#### 14 days after POST (6/10) PRE: 1.67 qt Bicep II Magnum *fb* POST: 1 fl oz Impact + 1 pt atrazine 4L + 1% v/v COC



©Dr. Rodrigo Werle, University of Wisconsin-Madison

#### 14 days after POST (6/10) PRE: 1.67 qt Bicep II Magnum *fb* POST: 3 fl oz Laudis + 1 pt atrazine 4L + 1% v/v COC



©Dr. Rodrigo Werle, University of Wisconsin-Madison